

Relating Linearly Superposed Periodic Solutions of Nonlinear Equations to One Soliton Solutions

W. Reinhardt^a, A. Khare^b, U. Sukhatme^c

^a Departments of Chemistry and Physics, University of Washington, Seattle, WA 98195-1700

^b Institute of Physics, Sachivalaya Marg, Bhubaneswar 751005, India

^c Department of Physics, State University of New York at Buffalo, Buffalo, NY 14260

Abstract. Using new generalized Landen transformations, we prove that the solutions of the KdV and other nonlinear equations obtained recently by using a kind of superposition principle for periodic solutions are in fact novel re-expressions of well known periodic one soliton solutions.

Nonlinear partial differential equations have a variety of unusual and sometimes unexpected solutions. For instance, it was not realized [1] until 1965 that the Korteweg-de Vries (KdV) equation [2]

$$u_t - 6uu_x + u_{xxx} = 0 , \quad (1)$$

supported multi-solitonic solutions, in addition to the single solitary wave solutions it was developed in 1895 to describe [3]. However, for traveling wave solutions of the form $u(x, t) = u(x - ct)$, the KdV equation has an integrating factor, and reduces to a nonlinear ordinary differential equation, immediately integrable to yield a Weierstrass function solution. In terms of the standard Jacobian elliptic functions $\text{sn}(x, m)$, $\text{cn}(x, m)$, $\text{dn}(x, m)$ [4] with argument x and modulus parameter m , a specific range of integration constants corresponds to the well-known cnoidal type periodic solution

$$u_1(x, t) = -2\alpha^2 \text{dn}^2[\alpha(x - b_1\alpha^2 t), m] + \beta\alpha^2 , \quad (2)$$

with velocity $b_1 = 8 - 4m - 6\beta$. It is worth noting that in the limit $m \rightarrow 1$, the period becomes infinite, and solution (2) reduces to the single soliton solution $-2\alpha^2 \text{sech}^2[\alpha(x - b_1\alpha^2 t)] + \beta\alpha^2$.

In a recent paper, Khare and Sukhatme [5] pointed out that certain kinds of linear superposition of the periodic solutions given in Eq. (2) are also solutions of the KdV equation by virtue of several new cyclic identities [6] satisfied by the Jacobi elliptic functions. The solutions are of the form

$$u_p(x, t) = -2\alpha^2 \sum_{i=1}^p d_i^2 + \beta\alpha^2 , \quad (3)$$

where p is any integer and

$$d_i \equiv \text{dn}[\alpha(x - b_p\alpha^2 t) + \frac{2(i-1)K(m)}{p}, m] . \quad (4)$$

The velocity is $b_p = 8 - 4m - 6\beta + 12A(p, m)$ with $A(p, m)$ being a constant in a certain cyclic identity [5].

In addition to the KdV equation, linear combination solutions similar to Eq. (3) were also shown to exist [7] for many other nonlinear differential equations. In the context of the static sine-Gordon equation, it was subsequently pointed out [8] that such linearly superposed solutions cannot be new, but must be re-expressible in terms of the initial periodic solution with different modulus parameter and integration constants. This connection was established by the discovery of new generalized Landen transformations [8, 9]. The point of this letter is to demonstrate that for the KdV equation also, the periodic solutions of the form (3), although they appear to be

new, can be simply re-expressed in terms of the original single Jacobi function solution given by Eq. (2).

Our procedure consists of starting with the recently found generalized Landen formula for $\text{dn}(x, \tilde{m})$ which involves a transformation from a modulus parameter m to a new modulus parameter $\tilde{m}$ [8, 9]:

$$\text{dn}(x, \tilde{m}) = \gamma [\text{dn}(\gamma x, m) + \text{dn}(\gamma x + 2K(m)/p, m) + \dots + \text{dn}(\gamma x + 2(p-1)K(m)/p, m)] , \quad (5)$$

where

$$\begin{aligned} \gamma &= [1 + \text{dn}(2K(m)/p, m) + \dots + \text{dn}(2(p-1)K(m)/p, m)]^{-1} , \\ \tilde{m} &= (m-2)\gamma^2 + 2\gamma^3[1 + \text{dn}^3(2K(m)/p, m) + \dots + \text{dn}^3(2(p-1)K(m)/p, m)] . \end{aligned} \quad (6)$$

Squaring both sides yields a Landen transformation for $\text{dn}^2(x, \tilde{m})$:

$$\text{dn}^2(x, \tilde{m}) = \gamma^2 \left[\sum_{i=1}^p \text{dn}^2[\gamma x + 2(i-1)K(m)/p, m] + \sum_{r=1}^p a_p(r) \right] , \quad (7)$$

where the constants $a_p(r)$ come from the cyclic identity $a_p(r) \equiv (d_1 d_{1+r} + \text{cyclic permutations})$ [6]. It now follows that solution (3) can be written as $-2\tilde{\alpha}^2 \text{dn}^2[\tilde{\alpha}(x - \tilde{c}t), \tilde{m}] + \tilde{\beta}\tilde{\alpha}^2$, provided one makes the identification

$$\tilde{\alpha} = \alpha/\gamma , \quad \tilde{c} = b_p \alpha^2 , \quad \tilde{\beta} = \beta \gamma^2 + 2\gamma^2 \sum_{r=1}^p a_p(r) . \quad (8)$$

Two other periodic traveling wave solutions of the KdV equation are

$$u_{\pm}(x, t) = \alpha^2 \left[m \text{sn}^2(\eta, m) \pm \sqrt{m} \text{cn}(\eta, m) \text{dn}(\eta, m) \right] , \quad \eta \equiv \alpha(x - q_1 \alpha t) , \quad (9)$$

with velocity $q_1 = -1 - m$. These solutions are different from Eq. (2) since they correspond to choosing a different range of integration constants in the general Weierstrass function solution of the KdV equation. They can also be obtained by making a Miura transform of the modified KdV equation. Linear superposition again gives apparently new solutions of the KdV equation, but in fact that is not the case. The solutions can be re-written in terms of the original solution (9) albeit with a change of parameters, using the generalized Landen transformation for $\text{sn}(x, \tilde{m})$ and $\text{cn}(x, \tilde{m})$ [8], with the same $\tilde{m}, m$ connection given in Eq. (6). A similar comment also applies to other nonlinear differential equations. Thus, although the idea of judicious linear superposition of periodic solutions has not given new solutions for nonlinear equations, it has certainly yielded

many novel, previously unexpected identities among the Jacobian elliptic functions themselves, some involving cyclic combinations [6] and others containing modulus parameter transformations [8, 9].

W.P.R. acknowledges the partial support of the US National Science Foundation, Grant PHY-0140091; and from NIST-NSF Digital Library of Mathematical Functions Project; U.S. thanks the U.S. Department of Energy for partial support of this research under grant DOE FG02-84ER40173.

References

- [1] N. J. Zabusky and M. D. Kruskal, Phys. Rev. Lett. **15** (1965) 240 .
- [2] D. J. Korteweg and G. de Vries, Phil. Mag.(5) **39** (1895) 422 .
- [3] P. G. Drazin and R. S. Johnson, *Solitons: An Introduction* (Cambridge Univ. Press, 1989).
- [4] See, for example, M. Abramowitz and I. Stegun, *Handbook of Mathematical Functions* (Dover, 1964).
- [5] A. Khare and U. P. Sukhatme, Phys. Rev. Lett. **88** (2002) 244101 .
- [6] A. Khare and U. P. Sukhatme, J. Math. Phys. **43** (2002) 3798 ; A. Khare, A. Lakshminarayan and U. P. Sukhatme, math-ph/0207019.
- [7] F. Cooper, A. Khare and U. P. Sukhatme, J. Phys. A: Math. Gen. **35** (2002) 10085.
- [8] A. Khare and U. P. Sukhatme, math-ph/0204054.
- [9] D. Lawden, *Elliptic Functions and Applications* (Springer-Verlag, 1989); A. Cayley, *An Elementary Treatise on Elliptic Functions* (Cambridge University Press, 1876).